

The planar Tree Lagrange Inversion Formula

Lothar Gerritzen

29.09.03

Introduction

A planar tree power series over a field K is a formal expression

$$\sum c_T \cdot T$$

where the sum is extended over all isomorphism classes of finite planar reduced rooted trees T and where the coefficients c_T are in K . Multiplications of these power series is induced by planar grafting of trees and turns the K -vectorspace $K\{x\}_\infty$ of those power series into an algebra, see [G].

If $f \in K\{x\}_\infty$ there is a unique $g(x) \in K\{x\}_\infty$ of order > 0 such that

$$g(x) = x \cdot f(g(x))$$

where $f(g(x))$ is obtained by substituting $g(x)$ for x in $f(x)$. Formulas for the coefficients of g in terms of the coefficients of f are obtained by the use of the planar tree Lukaciewicz language. This result generalizes the classical Lagrange inversion formula, see [C],[R],[Sch].

1 Lukaciewicz languages

Let Y be a $\mathbb{N}$ -graded set. It is given as a disjoint union of subsets Y_i for $i \in \mathbb{N}$. Assume that Y_0 is not empty.

Let $\mathbb{W}(Y)$ be the monoid of words over Y and let

$$\delta: \mathbb{W}(Y) \rightarrow \mathbb{N}$$

be the monoid morphism from $\mathbb{W}(Y)$ into additive monoid $\mathbb{N}$ of natural numbers for which

$$\delta(y) = i - 1$$

whenever $y \in Y_i$.

Definition 1.1. *Luk(Y) is the set of all words $w \in \mathbb{W}(Y)$ such that $\delta(w) = -1$ and such that if w' is a proper left factor of w , then $\delta(w') \geq 0$. We call Luk(Y) the Lukaciewicz language over the graded set Y .*

In case $\#Y_i = 1$ for all i , then $\text{Luk}(Y)$ is the classical Lukaciewicz language, for instance used in [C], (11.3). We will show that the results in [C], (11.3) remain correct in the more general situation.

Let $M(Y)$ be the submonoid of $\mathbb{W}(Y)$ generated by $\text{Luk}(Y)$.

Proposition 1.2. (i) *$M(Y)$ is a freely generated by $\text{Luk}(Y)$*

(ii) *Let $w \in \mathbb{W}(Y)$ with $\delta(w) = -r, r \in \mathbb{N}$, then $w \in M(Y)$ if and only if $r > 0$ and if w' is a proper left factor of w , then $\delta(w') > -r$.*

Proof. By definition $\text{Luk}(Y)$ has the prefix property and therefore $\text{Luk}(Y)$ is a code which means that $M(Y)$ is freely generated by $\text{Luk}(Y)$. $\square$

Example

$$\bigcup_{i=1}^{\infty} xi$$

then

$$\#Y_0 = 1$$

$$Y_0 \in Y_0$$

$$\mathbb{W}(Y+) \subseteq \text{Luk}$$

$$w \rightarrow wr + 1_0$$

If $w_1, \dots, w_m \in \text{Luk}(Y)$ and $y_m \in Y_m, m \geq 0$, then $v := y_m \cdot w_1 \cdot \dots \cdot w_m \in \text{Luk}(Y)$ Obviously

$$\begin{aligned} \delta(v) &= \delta(y_m) + \sum_{i=1}^m \delta(w_i) = \\ (m-1) + m(-1) &= -1. \end{aligned}$$

Let v' be a proper left factor of v , say $v = v' \cdot v''$. Then $v' = y_m w_1 \cdot \dots \cdot w_k \cdot w'_{k+1}$ with $0 \leq k < m$, w'_{k+1} is a proper factor of w_{k+1} .

Then

$$\begin{aligned} \delta(v') &= (m-1) + \sum_{i=1}^k \delta(w'_i) = \\ &= (m-k-1) + \delta(w'_{k+1}) \geq m-k-1 \geq 0 \end{aligned}$$

.

Proposition 1.3. *Let $w \in \text{Luk}(Y)$ and let $w(1)$ be the first letter of w . If $w(1) \in Y_m, m \geq 1$, then there are unique $w_1, \dots, w_m \in \text{Luk}(Y)$ such that*

$$w = w(1) \cdot w_1 \cdot w_2 \cdot \dots \cdot w_m$$

Proof. If $m = 0$, then $w = w(1)$, because otherwise $w(1)$ would be a proper left factor of w contained in $\text{Luk}(Y)$.

Let now $m > 0$. Then $\delta(w') = \delta(w) - \delta(w(1)) = (-1) - (m - 1) = -m$. There is a left factor w'_1 of w' of smallest length relative to Y such that

$$\delta(w'_1) = -1$$

Then $w'_1 \in \text{Luk}(Y)$. In a similar way $w'_2, \dots, w'_m \in \text{Luk}(Y)$ are determined. $\square$

Next we define the height $ht(v)$ of an element $v \in \text{Luk}(Y)$ by induction on the length of v .

If v is a letter in Y_0 , then $ht(v) := 0$.

Let now $v \in \text{Luk}(Y)$ and assume that the length $l_Y(v) = n > 1$. Then

$$v = v(1) \cdot v'_1 \cdot \dots \cdot v'_m$$

if

$$v(1) \in Y_m, m > 0.$$

and $l_Y(v'_i) < n$ for all i . By induction hypothesis we may assume that $ht(v'_i)$ is already defined for all i .

Let $ht(v) := 1 + \max_{i=1}^m ht(v'_i)$

Definition 1.4. *$ht(v)$ is called the height of $v \in \text{Luk}(Y)$.*

A classical result about Lukaciewicz languages is described now.

Let $\text{Luk}(\mathbb{N})$ be the Lukaciewicz language relative to the decomposition of $\mathbb{N}$ into subsets $\mathbb{N}_i = \{i\}$ for $i \in \mathbb{N}$. For $n \in \mathbb{N}$ we denote by $\lambda(n)$ the letter in $\mathbb{W} \in (\mathbb{N})$ represented by n in order that no confusion occurs between $\lambda(n) \cdot \lambda(m)$ and $\lambda(n \cdot m)$.

Let $\mathbf{PT}$ be the set of isomorphism classes of finite planar rooted trees, see [G 1] and $\mathbf{PT}' = \mathbf{PT} \cup \{1\}$ where 1 is the empty tree. The map

$$w : \mathbf{PT}' \rightarrow \text{Luk}(\mathbb{N})$$

is defined inductively with respect to the number $\#T^0$ of vertices of $T \in \mathbf{PT}'$.

If $T = 1$, let $w(T) := \lambda(0) \in \text{Luk}(\mathbb{N})$.

If $T = x$ is the tree in $\mathbf{PT}$ with a single vertex, then

$$w(x) = \lambda(1)\lambda(0) \in \text{Luk}(\mathbb{N}).$$

If $\#T^0 > 1$ and ρ_T is the root of T , then the subgraph $T - \rho_T$ obtained by deleting the vertex ρ_T and all the edges in T incident with ρ_T is a forest. Any component V_i of $T - \rho_T$ is a rooted tree whose root is the vertex in V_i incident with ρ_T in T . Also the components $V_1, \dots, V_m$, $m \geq 1$, are ordered and $\#V_i^0 < \#T$. We may assume that $w(V_1), \dots, w(V_m)$ are already defined. Let

$$w(T) := \lambda(m) \cdot w(V_1) \cdot \dots \cdot w(V_m).$$

Clearly $w(T) \in \text{Luk}(\mathbb{N})$

Proposition 1.5. *The map $w : \mathbf{PT} \rightarrow \text{Luk}(\mathbb{N})$ is bijective.*

Moreover $ht(w(T)) = \max_{b \in L(T)} (dist(\rho_T, b))$ where $dist(\rho_T, b)$ is the distance of the root ρ_T of T to a leaf b in T which is the length of a smallest path in T connecting ρ_T and b .

Proof. :1) Let $S, T \in \mathbf{PT}$ and assume that $w(S) = w(T)$.

Then

$$ar(S) = a(T) = m$$

If

$$m = 0$$

then $S = T$ is the empty tree.

If $m > 0$, then

$$T = T_1 \cdot \dots \cdot T_m$$

$$S = S_1 \cdot \dots \cdot S_m$$

and

$\#T_i^0 < \#T^0, \#S_i^0 < \#S^0$. If we proceed by induction on $\#T^0 + \#S^0$ we see that $T_i = S_i$ for all i . Thus $S = T$ and the map w is proved to be injective.

2) Let now $v \in \text{Luk}(\mathbf{PT})'$. If $v \in Y_0$, then $v = w(1)$, 1 empty tree. If the length of v is greater than 1,

than

$$v = v(1) \cdot v'_1 \cdot \dots \cdot v'_n$$

according to Proposition 1.4. Then $v(1) = \lambda(S), S \in \mathbf{PT}$ and

$$w(T_i) = v'_i$$

for


$$T_1, \dots, T_m \in \mathbf{PT}.$$

Then $deg(S) = m$ and $w(*_S(T_1, \dots, T_m)) = v$ where $*_S(T_1, \dots, T_m)$ is the planar grafting of $T_1, \dots, T_m$ over S . $\square$

Example 1.6. $v = \lambda(3)\lambda(0)\lambda(1)\lambda(2)\lambda(0)\lambda(1)\lambda(0) \in \text{Luk}(\mathbb{N})$

and $w(T) = v$ for

$T =$



2 The planar tree Lukaciewicz language

Let $\mathbf{PRT}$ be the set of isomorphism classes of finite planar reduced rooted trees and $\mathbf{PRT}' = \mathbf{PRT} \cup \{1\}$ where 1 denotes the empty tree, see [G1]. Let $\deg : \mathbf{PRT}' \rightarrow \mathbb{N}$ be the degree map defined by $\deg(1) = 0$, $\deg(T) = \#L(T)$ where $L(T)$ is the set of leaves of T . We denote by $\text{Luk}(\mathbf{PRT}')$ the Lukaciewicz language relative to the grading by $\deg$.

In the following we will give an interpretation of the elements in $\text{Luk}(\mathbf{PRT}')$ by right-sided decompositions of trees. For any tree $T \in \mathbf{PRT}$ and any vertex a of T , we denote by T_a the closed subtree in T generated by a .

There is a unique edge k in T incident with a such that T_a is a connected component of $T - k$ and such that the root ρ_T of T is the root in T_a . This gives another characterization of T_a .

Let $T \in \mathbf{PRT}$ and x be the tree in $\mathbf{PRT}$ with the single vertex.

Definition 2.1. T is called *right-sided*, if there is $T' \in \mathbf{PRT}'$ such that

$$T = x \cdot T'$$

The set of all right-sided trees in $\mathbf{PRT}$ is denoted by $x \cdot \mathbf{PRT}'$

Then

$$x = x \cdot 1, \quad x^2 = x \cdot x, \quad x \cdot x^2, \quad x \cdot x^3, \quad x \cdot (x^2 \cdot x), \quad x \cdot (x \cdot x^2) \in x \cdot \mathbf{PRT}'$$

Definition 2.2. S is called *completely right-sided* in T , if $\deg(S) > 1$ and for any leaf $b \in L(S)$ the closed subtree T_b in T generated by b is right-sided.

If S is completely right-sided in T , then the forest

$$T - \text{In}(S)$$

obtained by removing all the inner vertices of S from T is a disjoint union of right-sided trees.

Let S, S' be completely right-sided open subtrees of T and $S \subseteq S'$

Definition 2.3. *S is strictly contained in S' , if relative to T a leaf b of S which is also a leaf of S' is already a leaf of T .*

Let $T \in x \cdot \mathbf{PRT}$ and $r \geq 1$.

A right-sided open flag on T of length r , is a sequence

$$S = (S_1, S_2, \dots, S_r)$$

with:

$$S_r = T$$

S_i is a completely right-sided open subtree of T for all $i < r$

S_i is strictly contained in S_{i+1} for all $1 \leq i < r$.

Denote by $\Lambda_r(T)$ the set of all proper right-sided open flags on T .

Let $\Lambda_r = \cup \Lambda_r(T)$ where the union is extended over all trees in T for any $r \geq 1$.

Let

$$\Lambda = \bigcup_{r=0}^{\infty} \Lambda_r$$

with

$$\Lambda_0 = \{x\}$$

We are going to define a map

$$w : \Lambda \longrightarrow \text{Luk}(\mathbf{PRT}')$$

If $S \in \Lambda_0$, then $w(S) := \lambda(1) :=$ letter in $\mathbb{W}(\mathbf{PRT})'$ associated to the empty tree.

Let now

$$S \in \Lambda_r(T), r \geq 1, T \in x \cdot \mathbf{PRT}$$

If $r = 1$, then $S = T$ and $w(T) := \lambda(T') \cdot \lambda(1)^m$, if $T = x \cdot T'$, $\deg(T') = m$

Let now $r \geq 2$, $S = (S_1, \dots, S_r)$. We proceed by induction on $\deg(T)$.

Consider the planar forest

$$T - \text{In}(S_1) = V_0 \cup V_1 \cup \dots \cup V_m$$

and denote the connected components of $T - \text{In}(S_1)$ by $V_0, V_1, \dots, V_m$ Then $V_0 = x$ is the leaf of the left factor of T .

Let

$$S|V_i = (S_2 \cap V_i, \dots, S_r \cap V_i).$$

Then

$$S_r \cap V_i = V_i$$

and

$$S|V_i \in \Lambda_{r-1}(V_i)$$

for all i .

Define

$$w(S) := \lambda(S'_1) \cdot w(S|V_1)w(S|V_2) \cdot \dots \cdot w(S|V_m)$$

where

$$S_1 = x \cdot S'_1, \quad S'_1 \in \mathbf{PRT}.$$

As $w(S, V_i) \in \text{Luk } \mathbf{PRT}'$

and $\deg(S'_1) = m$ we get $w(S) \in \text{Luk}(\mathbf{PRT}')$

Theorem 2.4. *The map*

$$w : \Lambda \rightarrow \text{Luk}(\mathbf{PRT}')$$

is bijective.

3 Right-sided decompositions

Let $T \in \mathbf{PRT}$ and S be a non-empty subtree of T . The vertex in S closest to the root of T is defined to be the root ρ_S of S which turns S into a rooted tree.

Definition 3.1. *S is relatively open in T , if S is an open subtree in the closure $T_{(\rho_S)}$ of ρ_S in T .*

It is easy to show that S is relatively open in T if and only if $ar_S(a) = ar_T(a)$ for all vertices a of S with $ar_S(a) > 0$. There $ar_T(a)$ denotes the number of outgoing edges of the vertex a in T .

Let T be a right-sided tree in $\mathbf{PRT}$ of degree > 1 , $T = xT'$, and let Q be a system of relatively open subtrees U of T , $T = x \cdot T'$

Definition 3.2. *Q is called right-sided decomposition of T , if*

(i) *Each $U \in Q$ is right-sided*

(ii) $\bigcup_{U \in Q} U = T$

(iii) *If b is a leaf of T , then $\{b\} \in Q$ if and only if b is different from the first leaf of T .*

(iv) *If $U \cap U' \neq \emptyset$, $U \neq U'$, then $U \cap U' = \{a\}$, a vertex of T , and if a is not the root of U , then it is a leaf in T and the root of U' .*

Denote by $\Gamma(T)$ the set of all right-sided decompositions of T . Recall that $\Lambda(T)$ denotes the set of all proper open flags of T . For $S = (S_1, \dots, S_r) \in \Lambda(T)$ let $Q(S)$ denote the systems of subtrees of T which contains S_1 , all the connected components of $S_i - \text{In}(S_{i-1})$ for $1 < i \leq r$ and $\{b \in L(T) : b \neq \text{first leaf of } T\} := L(T')\}$, if $T = x \cdot T'$.

Then

$$Q(S) \in \Gamma(T)$$

Proposition 3.3. *The map*

$$S \longmapsto Q(S)$$

is a bijection


$$\Lambda(T) \rightarrow \Gamma(T)$$

for all

$$T \in x \cdot \mathbf{PRT}.$$

Example

$$T = x \cdot (x \cdot x^2 \cdot x \cdot x^3)$$



S = subgraph of T' vertices indexed by 1. S is open completely right-sided subtree of T and $T - \text{In}(S) = V_2 \cup \dots \cup V_5$ where V_i = subgraph of T of vertices indexed by i .

4 Lagrange inversion formula for planar tree power series

Let K be a field and $\mathbf{A} = K\{\{x\}\}_\infty$ the K -algebra of planar tree power series in x over K .

Proposition 4.1. *Let*

$$f \in \mathbf{A}, \text{ ord}(f) = 0$$

Then there is a unique power series

$$1/f \in \mathbf{A}$$

such that

$$f \cdot (1/f) = 1.$$

Then $\text{ord}(x \cdot (1/(f))) = 1$ and there is a unique $g \in \mathbf{A}$ such that

$$\text{ord}(g) = 1$$

$$\varphi_g(Sx1/f) = x$$

Proof. 1) The coefficients $\gamma(T)$ of $1/f$ satisfy the following system of equations:

$$c_1(f) \cdot \delta(1) = 1$$

$$c_x(f) \cdot \gamma(1) + c_1(f) \cdot \gamma(x) = 0$$

If

$$\text{ar}(T) = m \geq 2, T = T_1 \cdot T_2 \cdot \dots \cdot T_m$$

Then

$$c_T(F)\gamma(1) + c_1(f)\gamma(T) = 0$$

if

$$m > 2.$$

If

$$m = 2$$

Then

$$c_T(f)\gamma(1) + c_1(f) \cdot \gamma(T) + c_{T_1}(f)\gamma(T_2) = 0$$

There is a unique solution of this system of equation and $\sum \gamma(T) \cdot T$ is equal to $1/f$.

2) $\varphi_{(x \cdot 1/f)}$ is an automorphism of $\mathbf{A}$ as linear term of $x \cdot 1/f \neq 0$. Thus the inverse of $\varphi(x \cdot 1/(f))$ exists and is equal to φ_g for some $g \in \mathbf{A}$ Then φ_g for some $g \in \mathbf{A}$ Then

$$\varphi_g \circ \varphi_{(x \cdot 1/f)} = \varphi_x$$

and

$$\varphi_{g(x \cdot 1/f)} = x.$$

□

Proposition 4.2. Let $f \in \mathbf{A}$. Then there is a unique $g \in \mathbf{A}$ such that $g(x) = x \cdot f(g(x))$. It follows that $\text{ord}(g) \geq 1$.

Proof. 1) Let $a(t) = c_T(f)$ Now there is a unique system $b(T)$ of coefficients such that

$$b(T) = 0 \text{ if } T \neq x \cdot T'$$

$$b(x \cdot T') = \sum_{S' \in \Omega(T')} a(S')b(T' - \text{In}(S'))$$

Then

$$g = \sum b(T) \cdot T$$

satisfies the equation of the Proposition. 2) Let $b(T) = c_T(g)$. Then

$$c_T(x \cdot f(g(x))) = 0$$

if

$$T \neq x \cdot T'$$

and

$$\begin{aligned} c_{x \cdot T'}(x \cdot f(g(x))) &= c_{T'}(f(g(x))) \\ &= \sum_{S' \in \Omega(T')} b(T' - \text{In}(S')) \\ &= \sum_{\substack{S \in \Omega(T) \\ S \neq \delta_T}} a(S') b(T' - \text{In}(S)) \end{aligned}$$

Let

$$a(T) = c_T(f).$$

□

Corollary 4.3. :

$$g(x) = x \cdot f(g(x))$$

Proof.

$$\varphi_g(x \cdot 1/f) = g(x) \cdot (1/f)(g(x))$$

but

$$(1/f)(g(x)) = 1/f(g(x))$$

because

$$\text{ord}(f(g(x))) = 0.$$

□

Theorem 4.4. *Let f, g as in the Proposition above and*

$$a(T) = c_T(f), \quad b(T) = c_T(g).$$

Then

$$b(x) = a(1)$$

and if

$$T \in x \cdot \mathbf{PRT}, \quad \deg(T) > 1,$$

then

$$b(T) = \sum_{Q \in \Gamma(T)} a(Q)$$

where

$$a(Q) := \prod_{\substack{U \in Q \\ U=x \cdot U'}} a(U').$$

Proof. We proceed by induction on $n = \deg(T)$. If $n = 1$, it is clear. If $n > 1$, then

$$b(T) = \sum a(S) \cdot b(T - \text{In}(S))$$

where the summation is extended over all proper completely right-sided open subtrees S in T . By induction hypothesis

$$b(T - \text{In}(S)) = \prod_{i=1}^m b(V_i)$$

if

$$T - \text{In}(S) = V_1 \cup \dots \cup V_m$$

and

$$b(V_i) = \sum_{Q \in \Gamma(V_i)} a(Q)$$

The map $Q(T) \rightarrow Q(V_1) \times \dots \times Q(V_m)$ which maps S onto $(S(V_1), \dots, S(V_m))$ is a bijection. This proves the formula. $\square$

References

- [C] Cori, R.: Words and Trees, in M. Lothaire, Combinatorics on Words, Cambridge 1981, Chap 11, 213 - 227 Manuskript 2003
- [R] Raney, G. N. 1960: Functional composition patterns and power series reversion, Trans. Am. Math. Soc., 94, 441-451
- [Sch] Schützenberger, M. P. 1971: Le théorème de Lagrange selon Raney, in Logiques et Automates, Publications Inst. Rech. Informatique et Automatique, Rocquencourt, France.
- [G 1] Gerritzen, L.: Planar rooted trees and non-associative exponential series, Advances in Applied Mathematics 33 (2004) 342-365
- [G 2] Gerritzen, L.: Automorphisms of the planar tree power series algebra and the non-associative logarithm, Serdica Math. J. 30 (2004), 135-158
- [G 3] Gerritzen, L.: Planar Shuffle Product, co-Addition and the non-associative Exponential, ArXiv:math.RA/0502378